

## HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use PROSOL safely and effectively. See full prescribing information for PROSOL.

**PROSOL (amino acids) injection, for intravenous use**  
Initial U.S. Approval: 1998

### RECENT MAJOR CHANGES

Dosage and Administration, Preparation Instructions for Admixing Using a Parenteral Nutrition Container (2.3) 09/2020

### INDICATIONS AND USAGE

PROSOL is indicated as a source of amino acids for patients requiring parenteral nutrition when oral or enteral nutrition is not possible, insufficient, or contraindicated. PROSOL may be used to treat negative nitrogen balance in patents. (1)

### DOSAGE AND ADMINISTRATION

- *Pharmacy Bulk Package. Not for direct intravenous infusion.* (2.1)
- See full prescribing information for information on preparation, administration, instructions for use, dosing considerations, including the recommended dosage in adults and pediatrics, and dosage modifications in patients with renal impairment. (2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8)
- Protect the admixed parenteral nutrition solution from light. (2.3)

### DOSAGE FORMS AND STRENGTHS

Injection: 20% (0.2 grams/mL), 20 grams of amino acids per 100 mL in 2000 mL flexible containers. (3)

### CONTRAINDICATIONS

- Known hypersensitivity to one or more amino acids (4)
- Inborn errors of amino acid metabolism (4)
- Patients with pulmonary edema or acidosis due to low cardiac output (4)

### WARNINGS AND PRECAUTIONS

- Pulmonary Embolism due to Pulmonary Vascular Precipitates: if signs of pulmonary distress occur, stop the infusion and initiate a medical evaluation. (5.1)
- Hypersensitivity Reactions: monitor for signs and symptoms and discontinue infusion if reactions occur. (5.2)

- Risk of Infections, Refeeding Complications, and Hyperglycemia or Hyperosmolar Hyperglycemic State: monitor for signs and symptoms; monitor laboratory parameters. (5.3, 5.4, 5.5)
- Vein Damage and Thrombosis: solutions with osmolarity of 900 mOsmol/L or more must be infused through a central catheter. (2.2, 5.6)
- Hepatobiliary Disorders: monitor liver function parameters and ammonia levels. (5.7)
- Aluminum Toxicity: increased risk in patients with renal impairment, including preterm infants. (5.8, 8.4)
- Parenteral Nutrition Associated Liver Disease: increased risk in patients who receive parenteral nutrition for extended periods of time, especially preterm infants; monitor liver function tests, if abnormalities occur consider discontinuation or dosage reduction. (5.9, 8.4)
- Electrolyte Imbalance and Fluid Overload: patients with cardiac insufficiency or renal impairment may require adjustment of fluid, protein and electrolyte content. (5.10, 8.4)

### ADVERSE REACTIONS

Adverse reactions include pulmonary vascular emboli, hypersensitivity reactions infection, refeeding syndrome, hyperglycemia and hyperosmolar hyperglycemic state, vein thrombosis, elevated liver function tests, hyperammonemia, electrolyte imbalances, and hypervolemia. (6)

To report SUSPECTED ADVERSE REACTIONS, contact Baxter Healthcare Corporation at 1-866-888-2472 or FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).

### USE IN SPECIFIC POPULATIONS

Pediatric Use: increased risk of hypoglycemia/hyperglycemia: monitor serum glucose concentrations. (8.4)

See 17 for PATIENT COUNSELING INFORMATION

Revised: 10/2020

## FULL PRESCRIBING INFORMATION: CONTENTS\*

### 1 INDICATIONS AND USAGE

### 2 DOSAGE AND ADMINISTRATION

- 2.1 Important Preparation Information
- 2.2 Administration Instructions
- 2.3 Preparation Instructions for Admixing Using a Parenteral Nutrition Container
- 2.4 Dosing Considerations
- 2.5 Recommended Dosage in Adults
- 2.6 Dosage Modifications in Patients with Renal Impairment
- 2.7 Recommended Dosage in Pediatric Patients
- 2.8 Discontinuation of Parenteral Nutrition

### 3 DOSAGE FORMS AND STRENGTHS

### 4 CONTRAINDICATIONS

### 5 WARNINGS AND PRECAUTIONS

- 5.1 Pulmonary Embolism due to Pulmonary Vascular Precipitates
- 5.2 Hypersensitivity Reactions
- 5.3 Risk of Infections
- 5.4 Refeeding Syndrome
- 5.5 Hyperglycemia or Hyperosmolar Hyperglycemic State
- 5.6 Vein Damage and Thrombosis
- 5.7 Hepatobiliary Disorders

5.8 Aluminum Toxicity

5.9 Risk of Parenteral Nutrition Associated Liver Disease

5.10 Electrolyte Imbalance and Fluid Overload

5.11 Monitoring/Laboratory Tests

### 6 ADVERSE REACTIONS

### 8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

### 10 OVERDOSAGE

### 11 DESCRIPTION

### 12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action

### 15 REFERENCES

### 16 HOW SUPPLIED/STORAGE AND HANDLING

### 17 PATIENT COUNSELING INFORMATION

\* Sections or subsections omitted from the full prescribing information are not listed

## FULL PRESCRIBING INFORMATION

### 1 INDICATIONS AND USAGE

PROSOL is indicated as a source of amino acids for patients requiring parenteral nutrition when oral or enteral nutrition is not possible, insufficient, or contraindicated. PROSOL may be used to treat negative nitrogen balance in patients.

### 2 DOSAGE AND ADMINISTRATION

#### 2.1 Important Preparation Information

PROSOL is supplied as a pharmacy bulk package for *admixing only* and is *not for direct intravenous infusion*. Prior to administration, PROSOL *must be transferred to a separate parenteral nutrition container, diluted* and used as an admixture with or without dextrose, electrolytes and/or lipid emulsion.

- The key factor in preparation is careful aseptic technique to avoid inadvertent touch contamination during mixing of solutions and addition of other nutrients.
- Do not remove container from overpouch until ready to use.
- Tear protective overpouch across top at slit and remove solution container. Small amounts of moisture may be found on the solution container from water permeating from inside the container. The amount of permeated water is insufficient to affect the solution significantly. If larger amounts of water are found, the container should be checked for tears or leaks.
- Inspect PROSOL prior to use. Some opacity of the plastic due to moisture absorption during the sterilization process may be observed. This is normal and does not affect the solution quality or safety. The opacity will diminish gradually. Amino acid crystals may be present. Crystals will re-dissolve upon dilution during admixture compounding.

Evaluate the following:

- If the outlet port protector is damaged, detached, or not present, discard container as solution path sterility may be impaired.
- Check to ensure the solution is clear, colorless or slightly yellow. Discard if the solution is bright yellow or yellowish brown.
- Check for minute leaks by squeezing inner container. If leaks are found, discard container.
- PROSOL is intended for use in the preparation of sterile, intravenous admixtures. Because additives may be incompatible with PROSOL, evaluate all additions for compatibility.

#### 2.2 Administration Instructions

PROSOL is *for admixing use only*. It is *not for direct intravenous infusion*. Prior to administration, PROSOL *must be diluted* with other compatible intravenous fluids or used as an admixture with or without dextrose, electrolytes and/or lipid emulsion.

- PROSOL is to be used only in a suitable work area such as a laminar flow hood (or an equivalent

clean air compounding area). The key factor in the preparation is careful aseptic technique to avoid inadvertent touch contamination during mixing of solutions and addition of other nutrients.

- PROSOL is for admixing with dextrose injection and/or lipid emulsions using a parenteral nutrition container.
- When PROSOL is admixed with dextrose injection and/or lipid emulsion, the choice of a central or peripheral venous route should depend on the osmolarity of the final infusate. Solutions with greater than 5% dextrose or with osmolarity of 900 mOsmol/L or greater must be infused through a central venous catheter [see *Warnings and Precautions (5.6)*].
- Use a dedicated line for parenteral nutrition solutions.
- Intravenous lipid emulsions can be infused concurrently into the same vein as PROSOL-dextrose solutions by a Y-connector located near the infusion site; flow rates of each solution should be controlled separately by infusion pumps.
- For administration without lipid emulsion, use a 0.22 micron in-line filter. If lipid emulsion is also administered, use a 1.2 micron in-line filter.
- To prevent air embolism, use a non-vented infusion set or close the vent on a vented set, avoid multiple connections, do not connect flexible containers in series, fully evacuate residual gas in the container prior to administration, do not pressurize the flexible container to increase flow rates, and if administration is controlled by a pumping device, turn off pump before the container runs dry.
- If admixed or infused with lipid emulsion, do not use administration sets and lines that contain di-2-ethylhexyl phthalate (DEHP). Administration sets that contain polyvinyl chloride (PVC) components have DEHP as a plasticizer.
- Calcium and phosphate ratios must be considered. Excess addition of calcium and phosphate, especially in the form of mineral salts, may result in the formation of calcium phosphate precipitates [see *Warnings and Precautions (5.1)*].
- Prior to infusion, visually inspect the diluted parenteral nutrition solution containing PROSOL for particulate matter. The solution should be clear and there should be no precipitates. A slight yellow color does not alter the quality and efficacy of this product. If lipid has been added, ensure the emulsion has not separated. Separation of the emulsion can be visibly identified by a yellowish streaking or the accumulation of yellowish droplets in the mixed emulsion. Discard the admixture if any of the above are observed.

### **2.3 Preparation Instructions for Admixing Using a Parenteral Nutrition Container**

- Open by tearing protective overwrap across top at slit and remove solution container. If overpouch has been previously opened or is damaged, discard solution.
- If the outlet port protector is damaged, detached, or not present, discard the container.
- Some opacity of the plastic due to moisture absorption during the sterilization process may be observed.
- Check for minute leaks by squeezing the inner container firmly. If leaks are found, discard

solution.

- Once the protective foil overwrap has been removed, suspend container from eyelet support.
- Remove plastic protector from outlet port at bottom of container.
- Attach solution transfer set. Refer to complete directions accompanying set. Note: The closure shall be penetrated only one time with a suitable sterile transfer device or dispensing set which allows measured dispensing of the contents.
- Prepare the admixture into the parenteral nutrition container using strict aseptic techniques to avoid microbial contamination.
- Do not add the lipid emulsion to the parenteral nutrition container first; destabilization of the lipid emulsion may occur from such an admixture.
- PROSOL may be mixed with dextrose injection and/or lipid emulsion. The following proper mixing sequence must be followed to minimize pH related problems by ensuring that typically acidic dextrose injections are not mixed with lipid emulsions alone:
  1. Transfer dextrose injection to the parental nutrition container.
  2. Transfer PROSOL to the parenteral nutrition container.
  3. Transfer lipid emulsion to the parenteral nutrition container.
- Because additives may be incompatible, evaluate all additions to the parenteral nutrition container for compatibility and stability of the resulting preparation. Consult with pharmacist, if available. Questions about compatibility may be directed to Baxter. If it is deemed advisable to introduce additives to the parenteral nutrition container, use aseptic technique.
- Use gentle agitation during admixing to minimize localized concentration effects; shake containers gently after each addition.
- Alternatively, simultaneous transfer to the parenteral nutrition container of PROSOL, dextrose injection and lipid emulsion is also permitted.
- For admixing using an automated device, refer to Instructions for Use for the applicable device.
- The prime destabilizers of emulsions are excessive acidity (such as pH below 5) and inappropriate electrolyte content. Give careful consideration to additions of divalent cations ( $\text{Ca}^{++}$  and  $\text{Mg}^{++}$ ), which have been shown to cause emulsion instability. Amino acid solutions exert buffering effects that protect the emulsion.
- Inspect the final parenteral nutrition solution containing PROSOL to ensure that:
  - Precipitates have not formed during the mixing or addition of additives.

- The emulsion has not separated. Separation of the emulsion can be visibly identified by a yellowish streaking or the accumulation of yellowish droplets in the admixed emulsion.
- Discard the admixture if any of the above are observed.
- Containers should not be written on directly since ink migration has not been investigated. Affix accompanying label for date and time of entry.

### Stability and Storage

- Use for admixing should be limited to up to 4 hours at room temperature (25°C/77°F) after the container closure has been penetrated.
- Admixtures should be used promptly with storage under refrigeration 2° to 8°C (36° to 46°F) not to exceed 24 hours and must be completely used within 24 hours after removal from refrigeration. Any mixture remaining must be discarded.
- Protect the admixed parenteral nutrition solution from light.
- For single use only. Discard used container of PROSOL.

### **2.4 Dosing Considerations**

- PROSOL is for *admixing only*. *It is not for direct intravenous infusion*. Prior to administration, PROSOL *must be diluted* with other compatible intravenous fluids or used as an admixture with or without dextrose, electrolytes and lipid emulsion.
- PROSOL is a part of the parenteral nutrition regimen which also includes dextrose, electrolytes, and lipid emulsion. Protein, caloric, fluid and electrolyte requirements all need to be taken into consideration when determining individual patient dosage needs.
- The dosage of the final parenteral nutrition solution containing PROSOL must be based on the concentrations of all components in the solution and the recommended nutritional requirements [see *Dosage and Administration (2.5, 2.6, 2.7)*]. Consult the prescribing information of all added components to determine the recommended nutritional requirements for dextrose and lipid emulsion, as applicable.
- The dosage of PROSOL should be individualized based on the patient's clinical condition (ability to adequately metabolize amino acids), body weight and nutritional/fluid requirements, as well as additional energy given orally/enterally to the patient. Prior to initiating parenteral nutrition, the following patient information should be reviewed: review of all medications, gastrointestinal function and laboratory data (such as electrolytes (including magnesium, calcium, and phosphorus), glucose, urea/creatinine, liver panel, complete blood count and triglyceride level (if adding lipid emulsion)).
- Lipid emulsion administration should be considered with prolonged use (more than 5 days) of parenteral nutrition and is required in order to prevent essential fatty acid deficiency (EFAD). Serum lipids should be monitored for evidence of EFAD in patients maintained on fat-free parenteral nutrition. See complete prescribing information of lipid emulsion.

- Prior to administration of parenteral nutrition solution containing PROSOL, correct severe fluid, electrolyte and acid-base disorders.
- In many patients, provision of adequate calories in the form of hypertonic dextrose may require the administration of exogenous insulin to prevent hyperglycemia and glycosuria.
- Monitor levels of serum potassium during parenteral nutrition therapy. It may be necessary to add additional potassium to the parenteral nutrition admixture.

## 2.5 Recommended Dosage in Adults

The recommended adult daily dosage of PROSOL and the nutritional requirements for protein (nitrogen) are shown in **Table 1**.

As a component of parenteral nutrition, PROSOL provides 0.2 g protein/mL, which corresponds to 0.032 g nitrogen/mL.

As indicated on an individual basis, vitamins, electrolytes, trace elements and other components (including dextrose, electrolytes and lipid emulsion) can be added to the parenteral nutrition solution to meet nutrient needs and prevent deficiencies and complications from developing.

A maximum fluid supply of 40 mL/kg/day of parenteral nutrition solution, based on protein, should not be exceeded in adult patients; this volume does not take carbohydrates or electrolytes into consideration.

**Table 1. Recommended Daily Dosage in Adults**

Adult Patient Population	Recommended Protein <sup>a</sup> Requirement (g/kg/day) <sup>1</sup>	Corresponding Nitrogen Requirement (g/kg/day)	Recommended Daily Dosage of PROSOL (mL/kg/day) <sup>b</sup>
Stable Patients	0.8 to 1.0	0.13 to 0.16	4 to 5
Critically Ill Patients <sup>c</sup>	1.5 to 2.0	0.24 to 0.32	7.5 to 10

<sup>a</sup> Protein is provided as amino acids. When infused intravenously, amino acids are metabolized and utilized as the building blocks of protein.

<sup>b</sup> A maximum fluid supply of 40 mL/kg/day of parenteral nutrition solution should not be exceeded in adult patients.

<sup>c</sup> Includes patients requiring more than 2 to 3 days in the intensive care unit with organ failure, sepsis or postoperative major surgery. Do not use in patients with conditions that are contraindicated [see *Contraindications (4)*].

The flow rate of the parenteral nutrition solution must be adjusted taking into account the dose being administered, the daily volume intake, and the duration of the infusion. The flow rate should be increased gradually and governed, especially during the first few days of therapy, by the patient's tolerance to dextrose. Daily intake of PROSOL and dextrose should be increased gradually to the maximum required dose as indicated by frequent determinations of blood glucose levels.

## 2.6 Dosage Modifications in Patients with Renal Impairment

Prior to administration, correct severe fluid or electrolyte imbalances. Closely monitor serum electrolyte levels and adjust the volume of parenteral nutrition administered as required [see *Warnings and Precautions (5.10)*].

Patients with renal impairment not needing dialysis require 0.6 to 0.8 g of protein/kg/day. Serum electrolyte levels should be closely monitored. Patients on hemodialysis or continuous renal replacement therapy should receive 1.2 to 1.8 g of protein/kg/day up to a maximum of 2.5 g of protein/kg/day based on nutritional status and estimated protein losses.<sup>2</sup>

## 2.7 Recommended Dosage in Pediatric Patients

The dosage and constant infusion rate of intravenous dextrose must be selected with caution in pediatric patients, particularly neonates and low weight infants, because of the increased risk of hyperglycemia/hypoglycemia [see *Use in Specific Populations (8.4)*]. Frequent monitoring of serum glucose concentrations is required when dextrose is prescribed to pediatric patients, particularly neonates and low birth weight infants. The infusion rate and volume should be determined by the consulting physician experienced in pediatric intravenous fluid therapy.

In pediatric patients, PROSOL is dosed on the basis of protein provided as amino acids. The recommended dosage, by age group is provided in **Table 2**.

Infusion rates are based on protein and do not take carbohydrates, fluid or electrolytes into consideration.

PROSOL does not contain the amino acids cysteine and taurine, considered conditionally essential for neonates and infants. If possible, these amino acids should be added to the parenteral nutrition admixtures if used in this pediatric population.

**Table 2. Recommended Daily Dosage in Pediatric Patients**

Age	Recommended Protein <sup>a</sup> Requirement (g/kg/day) <sup>1</sup>	Corresponding Nitrogen Requirement (g/kg/day)	Recommended Daily Dosage of PROSOL (mL/kg/day)
Preterm and term infants less than 1 month of age	3 to 4	0.48 to 0.64	15 to 20
Pediatric patients > 1 month to 1 year of age	2 to 3	0.32 to 0.48	10 to 15
Pediatric patients 1 to 10 years of age	1 to 2	0.16 to 0.32	5 to 10
Pediatric patients 11 to 17 years of age	0.8 to 1.5	0.13 to 0.24	4 to 7.5

<sup>a</sup>Protein is provided as amino acids. When infused intravenously, amino acids are metabolized and utilized as the building blocks of protein.

## 2.8 Discontinuation of Parenteral Nutrition

To reduce the risk of hypoglycemia after discontinuation, a gradual decrease in flow rate in the last hour of infusion should be considered when administering parenteral nutrition solutions containing dextrose.

## 3 DOSAGE FORMS AND STRENGTHS

PROSOL 20% is a sterile solution of 20 grams of amino acids per 100 mL (0.2 gram/mL) available in 2000 mL flexible containers. **Table 3** describes the individual components of PROSOL.

**Table 3. Ingredients per 100 mL of PROSOL**

Amino Acids	20.0 g
Total Nitrogen	3.21 g
<b>Essential Amino Acids</b>	
Valine	1.44 g
Lysine (added as Lysine Acetate)	1.35 g
Histidine	1.18 g
Isoleucine	1.08 g
Leucine	1.08 g
Phenylalanine	1.00 g
Threonine	980 mg
Methionine	760 mg
Tryptophan	320 mg
<b>Nonessential Amino Acids</b>	
Alanine	2.76 g
Glycine	2.06 g
Arginine	1.96 g
Proline	1.34 g
Glutamic Acid	1.02 g
Serine	1.02 g
Aspartic Acid	600 mg
Tyrosine	50 mg
<b>Anion profiles per liter<sup>a</sup></b>	
Acetate from Lysine Acetate and glacial acetic acid <sup>b</sup>	140 mEq
pH (Range); pH adjusted with glacial acetic acid	6.0 (5.5 to 6.5)
<b>Osmolarity (calc)</b>	1835 mOsmol/L

<sup>a</sup> Balanced by ions from amino acids.

<sup>b</sup> Derived from pH adjustment with glacial acetic acid.

## 4 CONTRAINDICATIONS

The use of PROSOL is contraindicated in:

- Patients with known hypersensitivity to one or more amino acids [see *Warnings and Precautions (5.2)*].
- Patients with inborn errors of amino acid metabolism due to risk of severe metabolic or neurologic complications.
- Patients with pulmonary edema or acidosis due to low cardiac output.



## **5 WARNINGS AND PRECAUTIONS**

### **5.1 Pulmonary Embolism due to Pulmonary Vascular Precipitates**

Pulmonary vascular precipitates causing pulmonary vascular emboli and pulmonary distress have been reported in patients receiving parenteral nutrition. In some cases, fatal outcomes due to pulmonary embolism have occurred. PROSOL contains no added phosphorus. Patients, especially those with hypophosphatemia, may require the addition of phosphate. To prevent hypocalcemia, calcium supplementation should always accompany phosphate administration. Excessive addition of calcium and phosphate increases the risk of the formation of calcium phosphate precipitates. Precipitates have been reported even in the absence of phosphate salt in the solution. Precipitation following passage through an in-line filter and suspected in vivo precipitate formation has also been reported. If signs of pulmonary distress occur, stop the infusion and initiate a medical evaluation. In addition to inspection of the solution [see *Dosage and Administration* (2.1, 2.2, 2.3, 2.4)], the infusion set and catheter should also periodically be checked for precipitates.

### **5.2 Hypersensitivity Reactions**

Hypersensitivity reactions including anaphylaxis have been reported with parenteral nutrition solutions containing PROSOL. Stop the infusion immediately and treat patient accordingly if any signs or symptoms of a hypersensitivity reaction develop. Signs or symptoms may include: hypotension, hypertension, peripheral cyanosis, tachycardia, dyspnea, vomiting, nausea, urticaria, rash, pruritus, erythema, hyperhidrosis, pyrexia, and chills.

### **5.3 Risk of Infections**

Patients who require parenteral nutrition are at high risk of infections because the nutritional components of these solutions can support microbial growth. Infection and sepsis may also occur as a result of the use of intravenous catheters to administer parenteral nutrition.

The risk of infection is increased in patients with malnutrition-associated immunosuppression, hyperglycemia exacerbated by dextrose infusion, long-term use and poor maintenance of intravenous catheters, or immunosuppressive effects of other concomitant conditions, drugs, or other components of the parenteral formulation (e.g., lipid emulsion).

To decrease the risk of infection, ensure aseptic technique in catheter placement and maintenance, as well as aseptic technique in the preparation and administration of the nutritional formula.

Monitor for signs and symptoms (including fever and chills) of early infections, including laboratory test results (including leukocytosis and hyperglycemia) and frequent checks of the parenteral access device and insertion site for edema, redness and discharge.

### **5.4 Refeeding Syndrome**

Refeeding severely undernourished patients may result in refeeding syndrome, characterized by the intracellular shift of potassium, phosphorus, and magnesium as the patient becomes anabolic. Thiamine deficiency and fluid retention may also develop. To prevent these complications, monitor severely undernourished patients and slowly increase nutrient intakes.

## 5.5 Hyperglycemia or Hyperosmolar Hyperglycemic State

Administration of parenteral nutrition solutions containing dextrose in patients with diabetes mellitus, impaired glucose tolerance may worsen hyperglycemia. Administration of dextrose at a rate exceeding the patient's utilization rate may lead to hyperglycemia, coma, and death. Patients with underlying confusion and renal impairment who receive dextrose infusions, may be at greater risk of developing hyperosmolar hyperglycemic state. Monitor blood glucose levels and treat hyperglycemia to maintain optimum levels while administering parenteral nutrition solutions containing dextrose. Insulin may be administered or adjusted to maintain optimal blood glucose levels during administration of parenteral nutrition solutions containing dextrose.

## 5.6 Vein Damage and Thrombosis

PROSOL must be diluted and used as an admixture with or without dextrose, electrolytes and/or lipid emulsion. It is not for direct intravenous infusion. Solutions containing more than 5% dextrose or with an osmolarity of 900 mOsmol/L or greater must be infused through a central catheter [see *Dosage and Administration (2.2)*]. The infusion of hypertonic nutrient injections into a peripheral vein may result in vein irritation, vein damage, and/or thrombosis. The primary complication of peripheral access is venous thrombophlebitis, which manifests as pain, erythema, tenderness or a palpable cord. Remove the catheter as soon as possible, if thrombophlebitis develops.

## 5.7 Hepatobiliary Disorders

Hepatobiliary disorders are known to develop in some patients without preexisting liver disease who receive parenteral nutrition, including cholecystitis, cholelithiasis, cholestasis, hepatic steatosis, fibrosis and cirrhosis, possibly leading to hepatic failure. The etiology of these disorders is thought to be multifactorial and may differ between patients.

Increase in blood ammonia levels and hyperammonemia may occur in patients receiving amino acid solutions, including PROSOL. In some patients, this may indicate hepatic insufficiency or the presence of an inborn error of amino acid metabolism [see *Contraindications (4)*, *Use in Specific Populations (8.4)*].

Monitor liver function parameters and ammonia levels. Patients developing signs of hepatobiliary disorders should be assessed early by a clinician knowledgeable in liver diseases in order to identify possible causative and contributory factors, and possible therapeutic and prophylactic interventions.

## 5.8 Aluminum Toxicity

PROSOL contains no more than 25 mcg/L of aluminum. However, with prolonged parenteral administration in patients with renal impairment, the aluminum contained in PROSOL may reach toxic levels. Preterm infants are at a greater risk because their kidneys are immature, and they require large amounts of calcium and phosphate solutions, which contain aluminum.

Patients with renal impairment, including preterm infants, who receive parenteral levels of aluminum at greater than 4 to 5 mcg/kg/day, accumulate aluminum at levels associated with central nervous system and bone toxicity. Tissue loading may occur at even lower rates of administration.

## 5.9 Risk of Parenteral Nutrition Associated Liver Disease

Parenteral Nutrition Associated Liver Disease (PNALD) has been reported in patients who receive parenteral nutrition for extended periods of time, especially preterm infants, and can present as cholestasis or steatohepatitis. The exact etiology is unknown and is likely multifactorial. If patients

treated with parenteral nutrition solutions containing PROSOL develop liver test abnormalities, consider discontinuation or dosage reduction.

### **5.10 Electrolyte Imbalance and Fluid Overload**

Patients with renal impairment, such as pre-renal azotemia, renal obstruction, and protein-losing nephropathy may be at increased risk of electrolyte and fluid volume imbalance. Patients with cardiac insufficiency and pulmonary congestion are susceptible to excess fluid accumulation. Use parenteral nutrition solutions containing PROSOL with caution in patients with cardiac insufficiency or renal impairment. The dosage of parenteral nutrition may require adjustment with specific attention to fluid, protein, and electrolyte content in these patients.

Monitor renal function parameters. Patients developing signs of renal impairment should be assessed early by a clinician knowledgeable in renal disease in order to determine the appropriate parenteral nutrition dosage and other treatment options.

### **5.11 Monitoring/Laboratory Tests**

Monitor fluid and electrolyte status, serum osmolality, blood glucose, liver and kidney function, blood count and coagulation parameters throughout treatment. If electrolyte levels are severely elevated, stop parenteral nutrition containing PROSOL until levels have been corrected.

## **6 ADVERSE REACTIONS**

The following serious adverse reactions are discussed in greater detail in other sections of the prescribing information.

- Pulmonary embolism due to pulmonary vascular precipitates [see *Warnings and Precautions (5.1)*]
- Hypersensitivity reactions [see *Warnings and Precautions (5.2)*]
- Risk of infections [see *Warnings and Precautions (5.3)*]
- Refeeding syndrome [see *Warnings and Precautions (5.4)*]
- Hyperglycemia or hyperosmolar hyperglycemic state [see *Warnings and Precautions (5.5)*]
- Vein damage and thrombosis [see *Warnings and Precautions (5.6)*]
- Hepatobiliary disorders [see *Warnings and Precautions (5.7)*]
- Aluminum toxicity [see *Warnings and Precautions (5.8)*]
- Parenteral Nutrition Associated Liver Disease [see *Warnings and Precautions (5.9)*]
- Electrolyte imbalance and fluid overload [see *Warnings and Precautions (5.10)*]

The following adverse reactions from voluntary reports or clinical studies have been reported with parenteral amino acid products. Because many of these reactions were reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or establish a causal relationship to drug exposure.

- Metabolic acidosis
- Alkalosis
- Osmotic diuresis and dehydration
- Rebound hypoglycemia
- Hypo- and hyper-vitaminosis

## 8 USE IN SPECIFIC POPULATIONS

### 8.1 Pregnancy

#### Risk Summary

Limited published data with injectable amino acids solutions, including PROSOL in pregnant women are not sufficient to inform a drug associated risk for adverse developmental outcomes. However, malnutrition in pregnant women is associated with adverse maternal and fetal outcomes [see *Clinical Considerations*]. Animal reproduction studies have not been conducted with injectable amino acids solutions, including PROSOL.

The background risk of major birth defects and miscarriage for the indicated population is unknown. All pregnancies have a background risk of birth defect, loss, or other adverse outcomes. However, the background risk in the U.S. general population of major birth defects is 2 to 4% and of miscarriage is 15 to 20% of clinically recognized pregnancies.

#### Clinical Considerations

##### *Disease-Associated Maternal and/or Embryo-Fetal Risk*

Severe malnutrition in pregnant women is associated with preterm delivery, low birth weight, intrauterine growth restriction, congenital malformations and perinatal mortality. Parenteral nutrition should be considered if a pregnant woman's nutritional requirements cannot be fulfilled by oral or enteral intake.

### 8.2 Lactation

#### Risk Summary

There are no data available to assess the presence of injectable amino acids, including PROSOL in human milk, the effects of PROSOL on the breastfed infant or the effects on milk production. The lack of clinical data during lactation precludes a clear determination of the risk of PROSOL to a child during lactation; therefore, the developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for PROSOL and any potential adverse effects on the breastfed child from PROSOL or from the underlying maternal condition.

### 8.4 Pediatric Use

Neonates, especially premature infants with low birth weight, are at increased risk of developing hypo- or hyperglycemia and therefore need close monitoring during treatment with intravenous glucose solutions to ensure adequate glycemic control in order to avoid potential long term adverse effects [see *Dosage and Administration (2.7)*].

Plasma electrolyte concentrations should be closely monitored in the pediatric patients who may have impaired ability to regulate fluids and electrolytes.

Hyperammonemia is of special significance in infants (birth to two years). This reaction appears to be related to a deficiency of the urea cycle amino acids of genetic or product origin. It is essential that blood ammonia be measured frequently in infants [see *Warnings and Precautions (5.7)*].

Because of immature renal function, preterm infants receiving prolonged parenteral nutrition treatment with PROSOL may be at risk of aluminum toxicity [see *Warnings and Precautions (5.8)*].

Patients, including pediatric patients, may be at risk for PNALD [see *Warnings and Precautions (5.9)*].

### **8.5 Geriatric Use**

Clinical studies with PROSOL have not been performed to determine whether subjects aged 65 and over respond differently from other younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients.

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and of concomitant disease or drug therapy.

### **8.6 Renal Impairment**

In patients with impaired renal function, parenteral nutrition solutions containing PROSOL should be administered with caution. Frequent clinical evaluation and laboratory tests to monitor renal function such as serum electrolytes (especially phosphate and potassium) and fluid balance should be conducted [see *Dosage and Administration (2.6)*, *Warnings and Precautions (5.10)*].

### **8.7 Hepatic Impairment**

In patients with impaired liver function, parenteral nutrition solutions containing PROSOL should be administered starting at the low end of the dosing range [see *Dosage and Administration (2.5)*]. Frequent clinical evaluation and laboratory tests to monitor liver function such as bilirubin and liver function parameters should be conducted [see *Warnings and Precautions (5.7)*].

## **10 OVERDOSAGE**

An increased infusion rate of parenteral nutrition can cause hypervolemia, electrolyte disturbances, acidosis and/or azotemia, hyperglycemia, hyperosmolality [see *Warnings and Precautions (5.5, 5.10)*].

Severe hyperglycemia and severe dilutional hyponatremia, and their complications, can be fatal.

Discontinue infusion and institute appropriate corrective measures in the event of overhydration or solute overload during therapy, with particular attention to respiratory and cardiovascular systems.

For current information on the management of poisoning or overdosage, contact the National Poison Control Center at 1-800-222-1222 or [www.poison.org](http://www.poison.org).

## **11 DESCRIPTION**

PROSOL 20% (amino acids) injection is a sterile, nonpyrogenic, hypertonic solution of essential and nonessential amino acids supplied in a flexible container as a Pharmacy Bulk Package. A Pharmacy Bulk Package is a container of a sterile preparation for parenteral use that contains many single doses. The contents are intended for use in a pharmacy admixture program [see *Dosage and Administration (2.1)*]. PROSOL is not for direct intravenous infusion.

The formulas for the individual amino acids found in PROSOL are provided in **Table 4**.

**Table 4. Formulas for Amino Acids**

<b>Essential Amino Acids</b>	
Leucine	$(\text{CH}_3)_2 \text{CHCH}_2\text{CH}(\text{NH}_2) \text{COOH}$
Isoleucine	$\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3) \text{CH}(\text{NH}_2) \text{COOH}$
Lysine (added as the hydrochloride salt)	$\text{H}_2\text{N}(\text{CH}_2)_4 \text{CH}(\text{NH}_2) \text{COOH}$
Valine	$(\text{CH}_3)_2 \text{CHCH}(\text{NH}_2) \text{COOH}$
Phenylalanine	$(\text{C}_6\text{H}_5) \text{CH}_2 \text{CH}(\text{NH}_2) \text{COOH}$
Histidine	$(\text{C}_3\text{H}_3\text{N}_2) \text{CH}_2\text{CH}(\text{NH}_2) \text{COOH}$
Threonine	$\text{CH}_3\text{CH}(\text{OH}) \text{CH}(\text{NH}_2) \text{COO}$
Methionine	$\text{CH}_3\text{S}(\text{CH}_2)_2 \text{CH}(\text{NH}_2) \text{COOH}$
Tryptophan	$(\text{C}_8\text{H}_6\text{N}) \text{CH}_2 \text{CH}(\text{NH}_2) \text{COOH}$
<b>Nonessential Amino Acids</b>	
Alanine	$\text{CH}_3\text{CH}(\text{NH}_2) \text{COOH}$
Arginine	$\text{H}_2\text{NC}(\text{NH}) \text{NH}(\text{CH}_2)_3 \text{CH}(\text{NH}_2) \text{COOH}$
Aspartic Acid	$\text{HOOC} \text{CH}_2 \text{CH}(\text{NH}_2) \text{COOH}$
Glutamic Acid	$\text{HOOC}(\text{CH}_2)_2 \text{CH}(\text{NH}_2) \text{COOH}$
Glycine	$\text{H}_2\text{NCH}_2\text{COOH}$
Proline	$[(\text{CH}_2)_3 \text{NH} \text{CH}] \text{COOH}$
Serine	$\text{HOCH}_2\text{CH}(\text{NH}_2) \text{COOH}$
Tyrosine	$[\text{C}_6\text{H}_4(\text{OH})] \text{CH}_2\text{CH}(\text{NH}_2) \text{COOH}$

PROSOL contains no more than 25 mcg/L of aluminum.

## 12 CLINICAL PHARMACOLOGY

### 12.1 Mechanism of Action

PROSOL is used as a supplement of nutrition in patients, providing amino acids parenterally.

The amino acids provide the structural units that make up proteins and are used to synthesize proteins and other biomolecules or are oxidized to urea and carbon dioxide as a source of energy.

## 15 REFERENCES

1. Ayers P. et al. A.S.P.E.N. Parenteral Nutrition Handbook, 2nd ed. 2014 pg. 123
2. Mueller CM ed. The A.S.P.E.N. Nutrition Support Core Curriculum 3rd ed. 2017. Chapter 29 Sarav M, Kovesdy C. Renal Disease., pg. 565-586

## 16 HOW SUPPLIED/STORAGE AND HANDLING

PROSOL 20% (amino acids) injection is available in plastic Pharmacy Bulk Package flexible containers in the following size as shown in **Table 5** below.

**Table 5. PROSOL Dosage Form**

<b>Code</b>	<b>Volume</b>	<b>NDC Number</b>
2B6186	2000 mL	NDC 0338-0499-06

Minimize exposure of PROSOL to heat and avoid excessive heat.

Protect from freezing.

Store PROSOL at room temperature (25°C/77°F). Brief exposure up to 40°C/104°F does not adversely affect the product.

Do not use if protective overpouch has been previously opened or damaged.

For storage of admixed solutions [see *Dosage and Administration (2.3)*].

## **17 PATIENT COUNSELING INFORMATION**

Inform patients, caregivers, or home healthcare providers of the following risks of PROSOL:

- Pulmonary embolism due to pulmonary vascular precipitates [see *Warnings and Precautions (5.1)*]
- Hypersensitivity reactions [see *Warnings and Precautions (5.2)*]
- Risk of infections [see *Warnings and Precautions (5.3)*]
- Refeeding syndrome [see *Warnings and Precautions (5.4)*]
- Hyperglycemia or hyperosmolar hyperglycemic state [see *Warnings and Precautions (5.5)*]
- Vein damage and thrombosis [see *Warnings and Precautions (5.6)*]
- Hepatobiliary disorders [see *Warnings and Precautions (5.7)*]
- Aluminum toxicity [see *Warnings and Precautions (5.8)*]
- Parenteral Nutrition Associated Liver Disease (PNALD) [see *Warnings and Precautions (5.9)*]
- Electrolyte imbalance and fluid overload [see *Warnings and Precautions (5.10)*]

### **Baxter Healthcare Corporation**

Deerfield, IL 60015 USA

07-19-00-3506

Baxter and Prosol are registered trademarks of Baxter International Inc.